

LAMA3, LAMB3, COL17A1, AMBN, ENAM and WDR72 Mutations Causing AI

Objectives: Amelogenesis imperfecta (AI) can be manifested as isolated (non-syndromic) enamel malformations, or as a phenotype in syndromes. To date, 18 genes have been implicated in the etiology of isolated AI (*LAMA3*, *LAMB3*, *COL17A1*, *FAM83H*, *DLX3*, *AMELX*, *ENAM*, *AMBN*, *AMTN*, *C4orf26*, *ODAM*, *MMP20*, *KLK4*, *WDR72*, *ITGB6*, *SLC24A4*, *ACPT*, *GPR68*). In some cases, isolated AI is observed in the carriers of syndromes, as in the case of junctional epidermolysis bullosa (JEB). Syndromic AI patients are sometimes diagnosed as having isolated AI because the non-dental phenotypes go undetected or appear later in life. Genetic tests to determine the cause of AI can distinguish among isolated, carrier, or syndromic forms of AI, which improves patient management and prognosis. To identify the disease-causing mutations in families with AI.

Methods: AI families were analyzed by whole exome sequencing (WES) of genomic DNA from both parents and at least one affected offspring. Raw reads were aligned against human genome assembly GRCh37 using BWA. Variants were called and calibrated using GATK. High confidence variants were annotated and filtered by VarSeq. Potentially damaging mutations were prioritized, screened for each family member by Sanger sequencing to validate cosegregation of phenotype and genotype.

Results: The following AI-causing mutations were identified: heterozygous *LAMA3* (c.7367delG, p.Gly2456Alafs*22), heterozygous *LAMB3* (c.1705C>T, p.Arg569*), heterozygous *LAMB3* (c.3361G>T, p.Glu1121*), heterozygous *LAMB3* (c.3518G>T, p.*1173Leuext*56), *COL17A1* heterozygous (c.3327delT, p.Pro1110Argfs*21), compound heterozygous *AMBN* (c.1340C>T, p.Pro447Leu/c.1061T>C, p.Leu354Pro), *ENAM* heterozygous (c.588+1delG, splice junction), *ENAM* heterozygous (c.395dupA, p.Pro133Alafs*13), *WDR72* homozygous (c.377G>A, p.Trp126*), *WDR72* homozygous (c.1265G>T, p.Gly422Val) and *WDR72* homozygous c.1467_1468delAT, p.Val491fs*8).

Conclusions: We identified novel AI-causing mutations in multiple families with inherited enamel defects. Screening known AI candidate genes currently has a 40 to 50% chance of identifying the disease-causing mutation in a given family, suggesting that many more AI causing genes await discovery.

Division: AADR/CADR Annual Meeting
Meeting: 2018 AADR/CADR Annual Meeting (Fort Lauderdale, Florida)
Location: Fort Lauderdale, Florida
Year: 2018
Final Presentation ID: 1664

Authors

- Zhang, Hong (University of Michigan School of Dentistry , Ann Arbor Ann Arbor , Michigan , United States)
- Kasimoglu, Yelda (Faculty of Dentistry, Istanbul University , Istanbul Istanbul , Turkey)
- Koryuucu, Mine (Faculty of Dentistry, Istanbul University , Istanbul Istanbul , Turkey)
- Seymen, Figen (Faculty of Dentistry, Istanbul University , Istanbul Istanbul , Turkey)
- Kim, Jung-wook (School of Dentistry, Seoul National University , Seoul Seoul , Korea (the Republic of))
- Simmer, James (University of Michigan School of Dentistry , Ann Arbor Ann Arbor , Michigan , United States)
- Hu, Jan C.-c. (University of Michigan School of Dentistry , Ann Arbor Ann Arbor , Michigan , United States)

Support Funding Agency/Grant Number: NIDCR grant DE015846
Financial Interest Disclosure: NONE

SESSION INFORMATION

Poster Session
Mineralized Tissue IV
Saturday, 03/24/2018, 11:00AM - 12:15PM

[Back](#)

[Print Friendly](#)

