
ISSN 1306 - 696X

TURKISH JOURNAL of TRAUMA & EMERGENCY SURGERY

Ulusal Travma ve Acil Cerrahi Dergisi



11. ULUSAL TRAVMA ve ACİL CERRAHİ KONGRESİ

11TH NATIONAL CONGRESS ON TRAUMA AND EMERGENCY SURGERY

April 5-9, 2017 Cornelia Diamond Hotel, Belek, Antalya

BİLDİRİ ÖZETLERİ / ABSTRACTS

Volume 23 | Supp. 1 | April 2017

www.tjtes.org

SB-067

Paraözofageal Herniye Bağlı Sıradışı Proksimal Gastrik Nekroz: Olgu Sunumu

ÖMER CENK CÜCÜK, ADEM BAYRAKTAR, MEHMET İLHAN, MEVLÜT YORDANAGİL, FİRUZ GACHAYEV, RECEP ERÇİN SÖNMEZ, CEMALETTİN ERTEKİN

İstanbul Üniversitesi İstanbul Tıp Fakültesi, Genel Cerrahi Anabilim Dalı, İstanbul

Giriş: Paraözofageal herni (Tip II diyafram hernisi), mide ve diğer karın içi organların peritoneal fıtık kesesi ile birlikte hiatustan posterior mediastene doğru yer değiştirmesidir. Paraözofageal herniler, etyolojisi belli olmayan ve sonradan gelişen bir fıtık şeklidir. En önemli komplikasyonu fıtık kesesi içindeki organların strangülasyonu ve perforasyonudur. Bu komplikasyonlar hastaların üçte birinde görülür ve acil cerrahi müdahale gerekir. Karın ağrısı ile acil cerrahi polikliniğine başvuran ve yapılan tetkiklerinde paraözofageal herniye sekonder midede strangülasyon saptanan ve proksimal gastrektomi+gastroözofagostomi yapılan hastayı sunmayı amaçladık. **Olgu:** Karın-göğüs ağrısı, iştahsızlık, bulantı ve öğürme şikayetleri ile acil cerrahi polikliniğine başvuran 78 yaşında hipertansiyon dışında ek hastalığı olmayan kadın hasta. Fizik muayenede epigastrik bölgede daha fazla olmak üzere batında yaygın hassasiyet saptandı. Laboratuar tetkiklerinde lökositoz (14500/μL) ve akut faz reaktanlarının yükseldiği (CRP: 9 mg/L) gözlendi. Akciğer grafisinde sol diyaframın üstünde içinde oral kontrast olan mide ile uyumlu görünüm saptandı. Çekilen oral ve intravenöz kontrastlı batın bilgisayarlı tomografide (BT) midenin fundus ve korpusunun büyük kısmının sol hemitoraks içerisinde olduğu görüldü. Preoperatif kardiyak arrest sonrası yapılan kardiyopulmoner resüsistasyona cevap alındı ve laparotomide midenin paraözofageal herniye bağlı fundusunun ve korpusunun büyük kısmının toraksa geçtiği ve sıkışmaya bağlı nekroz ve perforasyon geliştiği gözlendi. Proksimal gastrektomi ve özofagogastrik anastomoz yapıldı. Perforasyona bağlı kontaminasyon ve hastanın genel durum bozukluğu nedeniyle diyaframa mesh takviyesi ile tamir yapılmadı. Postoperatif 5. günde oral sulu gıda başlandı. Takiplerinde vitalleri ve genel durumu stabil seyreden hasta önerilerle taburcu edildi. **Tartışma ve Sonuç:** Paraözofageal herniler nadir görülen diyafram hernilerindendir ve organ strangülasyonu ciddi ve hayatı tehdit eden komplikasyonudur. Bu olgularda cerrahi tedavi şart ve hayat kurtarıcıdır.

Unusual Proximal Gastric Necrosis Due to Paraoesophageal Hernia: Case Report

ÖMER CENK CÜCÜK, ADEM BAYRAKTAR, MEHMET İLHAN, MEVLÜT YORDANAGİL, FİRUZ GACHAYEV, RECEP ERÇİN SÖNMEZ, CEMALETTİN ERTEKİN

Department of General Surgery, İstanbul University İstanbul Faculty of Medicine, İstanbul

Introduction: Paraoesophageal hernia (Type II diaphragm hernia) is the displacement of the stomach and other abdominal organs along with the peritoneal sac to the posterior mediastinum. Paraoesophageal hernia is a herniation form that has no definite etiology and develops at late ages. The most important complication is strangulation and perforation of the organs within the hernia sac. These complications are seen in one third of the patients and urgent surgical intervention is needed. We aimed to present a patient who admitted to emergency department with abdominal pain and had gastric strangulation secondary to paraoesophageal hernia and underwent proximal gastrectomy+g astrooesophagostomy. **Case:** A 78-year-old woman who have no additional disease except hypertension, was admitted to the emergency surgery clinic with abdominal-chest pain, anorexia, nausea and retching. Physical examination revealed widespread tenderness in the abdomen, more in the epigastric region. Leukocytosis (14500/μL) and elevated acute phase reactants (CRP: 9 mg/L) were observed in initial laboratory tests. On chest X-ray, a stomach-like appearance with oral contrast was found on the left diaphragm. Oral and intravenous contrasted abdominal computed tomography (CT) revealed that majority of the stomach were in the left hemithorax. After preoperative cardiac arrest, cardiopulmonary resuscitation was applied and in laparotomy majority of the fundus and corpus of the stomach was displaced to thorax, necrosis and perforation on gastric wall was identified due to compression. Proximal gastrectomy and oesophagogastric anastomosis were performed. Due to perforation-related contamination and deterioration in the general condition of the patient, diaphragm repair was not performed with mesh reinforcement. On the fifth postoperative day oral food intake was started. Patient with stable vital signs and general condition was discharged with the suggestions. **Discussion and Conclusion:** Paraoesophageal hernia is a rare type of diaphragmatic hernias and organ strangulation is a serious and life-threatening complication. Surgical treatment is a condition and life saver in these cases.

SB-068

Akut Divertikülit Tanısıyla Tedavi Edilen Hastaların Sonuçlarının Değerlendirilmesi

GÖRKEM YILDIZ, AMAN GAPBAROV, İHSAN METİN LEBLEBİCİ, NURAY ÇOLAPKULU, TUĞRUL ÖZDEMİR, ŞÜKRİYE JÜLİDE SAĞIROĞLU, ÖZGÜR EKİNCİ, GÜRHAN BAŞ, ORHAN ALİMOĞLU

İstanbul Medeniyet Üniversitesi Tıp Fakültesi Göztepe Eğitim ve Araştırma Hastanesi, Genel Cerrahi Anabilim Dalı, İstanbul

Amaç: Kolon divertikülleri toplumumuzda zamanla sıklığı artsa da halen nadir görülmektedir. Bu çalışmada akut divertikülit tanısı ile hastanemiz genel cerrahi servisine yatırılıp tedavi edilen hastalarımızı ve sonuçlarını sunmayı amaçladık. **Gereç ve Yöntem:** 01 Ocak 2016 ve 31 Aralık 2016 tarihleri arasında akut divertikülit tanısı ile tedavi edilen hastalar retrospektif olarak incelendi. Hastalar yaş, cinsiyet, uygulanan tedavi, komplikasyon ve rekürren atak geçirmesine göre değerlendirildi. **Bulgular:**

SB-068

Evaluation of Outcomes in Patients Treated for Acute Diverticulitis

GÖRKEM YILDIZ, AMAN GAPBAROV, İHSAN METİN LEBLEBİCİ, NURAY ÇOLAPKULU, TUĞRUL ÖZDEMİR, ŞÜKRİYE JÜLİDE SAĞIROĞLU, ÖZGÜR EKİNCİ, GÜRHAN BAŞ, ORHAN ALİMOĞLU

Department of General Surgery, İstanbul Medeniyet University Faculty of Medicine Göztepe Training and Research Hospital, İstanbul

Objective: Colonic diverticula are still rare even though incidence increase over time. We aimed to present the outcome of the patients treated for acute diverticulitis in general surgery clinic. **Material and Method:** Patients treated for acute diverticulitis between 01 February 2016 and 31 December 2016 were examined retrospectively. The patients were evaluated according to their age, sex, treatment method, recurrence and complications. **Findings:** Thirty-six patient were trea-